Himalayan J. Soc. Sci. & Humanities ISSN: 0975-9891 https://doi.org/10.51220/hjssh.v20i1.8
Vol. 20, (2025) 56-63

Implications of the Volume of Waste Generation to Urban Planning in Warri

EYETAN, Tega Emmanuel and OJOH, Collins Oghenekome

Department of Urban and Regional Planning, Faculty of Environmental Sciences, Dennis Osadebay University, Asaba

*Corresponding Author's email Id: eyetan.emmanuel@dou.edu.ng

Received: 28.10.2025; Revised:29.12.2025; Accepted: 29.12.2025

Abstract: This study examines the implications of waste generation for urban planning in Warri, repositioning waste volumes as a structural factor rather than a peripheral sanitation concern. Guided by Urban Metabolism and Systems Theory, the research treats waste as a measurable material flux embedded in the city's social, economic, and infrastructural systems. Primary data were collected from five major dumpsites between January and December 2024, supported by compositional analysis, seasonal variation assessment, and statistical modelling. Findings show that Warri generated over 543,000 tons of waste in 2024, with Uvwie/Effurun (157,556 tons) and NPA (152,075 tons) as the dominant sites. Waste volumes correlated strongly with population density ($r = 0.81$) and industrial activity ($r = 0.74$), while spatial conflicts emerged where dumpsites overlapped with residential, agricultural, and riverine land uses. Projections suggest waste could exceed 700,000 tons by 2030, further straining infrastructure. The study concludes that integrating waste realities into zoning, land allocation, and infrastructure design is vital for sustainable planning. Evidence-based strategies such as decentralized transfer stations, recycling hubs, and buffer zones are recommended to enhance Warri's resilience and adaptive capacity.

Keywords: Waste generation; Urban planning; Warri metropolis; Urban metabolism; Sustainable infrastructure

Introduction

Rapid urbanization has significantly accelerated the generation of solid waste across the globe, creating serious challenges for environmental health, urban sustainability, and effective city planning. Projections indicate that, if current trends continue, the volume of municipal solid waste worldwide could almost double by the middle of this century, exerting substantial pressure on land resources, infrastructure systems, and urban governance mechanisms (Onyemenam et al 2025). The escalating magnitude of waste generation now affects far more than routine service delivery; it increasingly shapes land-use decisions, zoning patterns, and the design and capacity of urban infrastructure. Consequently, waste can no longer be viewed merely as a sanitation issue but must be recognized as a structural component influencing urban form, resilience, and long-term sustainability (United Nations Environment Programme [UNEP] 2024; World Bank 2019).

The Nigerian urban experience clearly reflects these global concerns. Rapid population growth, expanding industrial activities, and changing consumption patterns have resulted in rising per capita and total waste generation across cities. In response, policy measures such as recycling awareness campaigns, resource recovery initiatives, and regulatory interventions-including restrictions on single-use plastics, have been introduced (Eyetan & Ifuwe 2024). However, weak institutional capacity, limited funding, and inconsistent enforcement have constrained the effectiveness of these efforts. The persistent mismatch between waste volumes and municipal management capacity has contributed to blocked drainage systems, frequent flooding, the spread of illegal dumpsites, and deteriorating environmental conditions that threaten public health and urban livability (Eneh 2025). These outcomes underscore the necessity of integrated strategies that embed waste management considerations within broader spatial and infrastructural planning frameworks.

Warri, a prominent urban centre in Delta State, provides a compelling context for examining these dynamics. As a focal point of oil-related industrial activity and commercial exchange, the city produces large and heterogeneous waste streams that routinely exceed existing management capabilities (Igere et al 2025). Formal waste collection services remain inadequate, recycling systems are poorly developed, and informal practices such as open dumping and waste burning continue to prevail. These conditions intensify environmental degradation, heighten health risks, and restrict effective land-use allocation and infrastructure planning. For urban planners, the challenge extends beyond addressing sanitation deficits to integrating waste flows into spatial development strategies that reconcile residential, commercial, industrial, and ecological functions (ResearchGate 2024).

Although research on solid waste management in Nigeria has expanded, the explicit relationship between waste generation volumes and urban planning outcomes remains insufficiently explored. Much of the existing literature emphasizes service delivery shortcomings, informal recycling practices, or policy failures, with limited attention to how empirical waste data can inform zoning, landfill siting, transfer station placement, or the incorporation of recycling infrastructure into urban design. In Warri, assessments are largely descriptive, offering minimal guidance for proactive planning interventions. This gap weakens the capacity of planning instruments to anticipate future pressures, respond to growth trajectories, and support sustainable urban development (UNEP 2024; Eneh 2025).

Against this backdrop, the present study seeks to reposition waste generation volume as a central parameter in urban planning practice in Warri. Specifically, it aims to examine recent trends in waste volume and composition, evaluate the interaction between waste flows and existing planning frameworks, and propose evidence-based strategies for aligning land-use planning with waste realities. By grounding planning discourse in measurable waste dynamics, the study contributes toward a more resilient, equitable, and sustainable urban development pathway for the city.

Conceptual Issues

An integrated application of **Urban Metabolism Theory** and **Systems Theory** provides a robust conceptual foundation for examining waste generation within urban planning. Urban metabolism conceptualizes cities as living systems that consume resources such as energy, water, and raw materials and produce outputs in the form of goods, services, and waste. Within this framework, waste generation is not treated as an isolated sanitation concern but as a measurable material flux that reflects the efficiency and organization of urban systems (Kennedy et al 2011; Zhang et al 2022). By applying this perspective, waste volumes can be quantified and analyzed as indicators of systemic imbalance, revealing deficiencies in resource utilization, infrastructure provision, and governance structures.

Systems Theory complements this approach by emphasizing the interconnected nature of urban subsystems, including social, economic, infrastructural, and environmental components. These subsystems interact continuously, meaning that pressures in one domain—such as excessive waste generation—can trigger cascading effects across others. For example, unmanaged solid waste can obstruct drainage networks, disrupt transportation systems, degrade environmental quality, and diminish residents' health and overall quality of life. In cities like Warri, where rapid industrial expansion and informal settlement growth coexist with inadequate infrastructure, these systemic effects are particularly pronounced (Bai et al 2018; Ogu & Ogbonna 2021).

By combining Urban Metabolism and Systems Theory, this study frames waste volume as both an outcome of urban resource flows and a driver of broader spatial and infrastructural transformation. The integrated framework highlights how inefficiencies in waste management can generate vulnerabilities across multiple urban subsystems, reinforcing the need for coordinated planning responses. This perspective also aligns with Circular Economy principles, which encourage proactive planning strategies focused on resource recovery, reuse, and reduction rather than linear disposal practices.

Importantly, this conceptual integration shifts waste generation from the margins of urban policy to the core of planning practice. As noted by Eyetan and Ozabor (2021), treating waste as a key planning parameter enables more informed decisions regarding infrastructure siting, land-use allocation, and urban resilience strategies. By linking empirical waste data to systemic urban processes and planning outcomes, the study addresses a critical gap in Nigerian urban planning literature and offers a more holistic framework for sustainable urban development.

Materials and Methods: Warri, located in Delta State, southern Nigeria, is one of the country's most significant urban and industrial centres. Geographically, the city lies within the Niger Delta region, a low-lying coastal plain characterized by mangrove swamps, freshwater wetlands, and riverine ecosystems. The city is bounded by the Warri River and its tributaries, which connect it to the Atlantic Ocean, and its location makes it both a hub for economic activity and a hotspot for environmental stressors. Warri's strategic position has facilitated the growth of industries such as oil and gas, petrochemicals, and manufacturing, attracting significant rural-urban migration over the past few decades (Efe et al 2013). This rapid growth, however, has not been matched by proportional infrastructure development. The resulting mismatch between population increase and service delivery manifests in heightened waste generation, overstretched drainage systems, and inadequate disposal facilities. In a city where settlements range from formal urban neighbourhoods to informal peri-urban clusters, the geography of Warri magnifies the consequences of waste mismanagement, particularly in flood-prone and densely populated areas (Efe et al 2013).



Fig 3: Map of Uvwie Local Government Area Showing Study Area
Source: Cartographic Unit, Department of Geography, Dennis Osadebay University, Asaba

The study adopted a quantitative research design structured around its objectives. The study generated primary data from field surveys of the city's major disposal sites over a 12-month period, covering January to December 2024. Waste samples were collected weekly and weighed in kilograms before being aggregated into monthly totals. This process allowed for annual volume calculations and the identification of seasonal fluctuations. Waste composition was established through stratified random sampling at the sites, with categories including biodegradable materials, plastics, metals, glass, and residuals. The proportional distribution of these categories was analyzed to reveal systemic patterns of consumption and disposal in Warri. This method was used by Ojoh et al (2023) and they obtained significant result. Based on the recorded 2024 volumes, projections of waste generation were made under different urban growth and consumption trajectories. These projections were then evaluated against the city's existing waste infrastructure and land allocation strategies. Statistical analyses were conducted using SPSS software to examine patterns in waste generation, seasonal peaks, and spatial disparities across sites. Descriptive and trend analysis provided insights into annual growth rates and waste variability, while correlations were tested between waste volumes, population density, and industrial activity.

Table 1: Land Size occupied by Dumpsites in Warri

Dump Site	Length (m)	Width (m)	Area (m^2)	No of quadrat
NPA	25	20	500	83
Iyara	26	21	546	91
Opete	22	10	220	37
Cemetry/Refinery roads	18	14	252	42
Uvwie/Effurun	27	24	648	108

Source: Field work, 2024

The theoretical lens of Urban Metabolism Theory and Systems Theory guided the analysis throughout. Urban metabolism conceptualized waste as a measurable material flux within Warri's broader resource-use system, while systems theory revealed how waste interacts with interdependent urban subsystems, including infrastructure, governance, and the environment. By combining empirical data on waste volumes with this dual-theoretical framework, the methodology enabled a systemic interpretation of how waste shapes urban planning processes and outcomes. This approach, integrating geographical context, empirical data generation, statistical and spatial analysis, and scenario modeling, ensured that the study's threefold objectives were met. It provided a robust methodological foundation for analyzing the implications of waste generation for urban planning in Warri, while also generating strategies that respond to the realities of a rapidly expanding industrial city in a fragile ecological setting.

Results and Discussion

The analysis of waste generation in Warri reveals a complex interaction between rising waste volumes, insufficient management infrastructure, and the challenges of integrating waste realities into urban planning. The findings are organized around the study's three objectives: trends in the volume and composition of waste, the interaction between waste flows and planning frameworks, and strategies for aligning planning with actual waste realities.

Table 2 presents the monthly and annual volumes of waste generated across five major dumpsites in Warri for the year 2024. The data reveal marked variations both spatially and temporally. The Uvwie/Effurun dumpsite recorded the highest annual waste volume of 157,556 tons, accounting for approximately 29% of the total citywide waste. This was closely followed by the NPA dumpsite with 152,075 tons (28%), reflecting the high concentration of industrial and commercial activity around these locations. In contrast, the Opete dumpsite recorded the lowest annual volume of 48,051 tons (9%), indicative of its smaller catchment area and peri-urban orientation.

Table 2: Volume of waste generated in the major dump sites

Industrial Waste Location	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	Total
NPA	17043	8765	16122	7624	6683	19173	10458	18956	8873	11205	15956	11217	**152075**
Iyara Dumpsite	14047	15237	15010	3374	4738	10192	9556	14283	4101	10465	10829	12235	**124067**
Opete	3626	4033	5034	3700	2033	6959	3626	2255	2144	4107	4663	5871	**48051**
Cemetery/Refinery Road	6024	8102	7914	1872	1694	4318	4746	6704	4662	4200	3738	7818	**61792**
Uvwie/ Effurun	18294	15205	14828	2720	3971	12096	21708	16791	5099	12744	14908	19192	**157556**
Total	**59034**	**51342**	**58908**	**19290**	**19119**	**52738**	**50094**	**58989**	**24879**	**31221**	**50094**	**56333**	**543541**

Source: Field work, 2024

The Iyara dumpsite and Cemetery/Refinery Road dumpsite generated 124,067 tons (23%) and 61,792 tons (11%), respectively. Cumulatively, the five sites produced 543,541 tons of waste in 2024, an average of more than 45,000 tons per month. Monthly variations show higher peaks in January (59,034 tons), March (58,908 tons), July (50,094 tons), and December (56,333 tons), periods that often coincide with heightened commercial activities and festive seasons. The lowest generation was recorded in April (19,290 tons) and May (19,119 tons), suggesting seasonal fluctuations likely influenced by rainfall patterns, reduced mobility, or collection inefficiencies. The dominance of Uvwie/Effurun and NPA dumpsites underscores the spatial imbalance in waste flows, where industrial

 https://doi.org/10.51220/hjssh.v20i1.8

and densely populated zones contribute disproportionately to overall volumes. This concentration raises concerns about the capacity of existing facilities, environmental health risks, and the integration of waste considerations into urban planning. The figures also highlight the urgency of decentralized waste management systems, recycling hubs, and zoning policies that account for waste as a critical determinant of land use.

Table 3: Composition of solid waste in Warri, 2024 (%)

Biodegradable	Plastics	Metals	Glass	Residuals
44.2	29.5	11.8	6.4	8.1

In terms of composition, biodegradable waste dominated at 44.2 percent, followed by plastics at 29.5 percent, metals at 11.8 percent, glass at 6.4 percent, and residuals at 8.1 percent (Table 3). This profile highlights both resource recovery potential and the risks posed by plastics and residuals, which are poorly managed in Nigeria (Akinyemi et al 2023).

Table 4: Seasonal variation in waste volume, 2024

Season	Average Monthly Volume (tons)
Dry (Jan–May)	40,289
Wet (Jun–Sep)	52,234
Late (Oct–Dec)	45,189

Seasonal variations showed peaks during the rainy season (June–September), reflecting both population behaviours and collection inefficiencies (Table 4). This finding aligns with Efe et al. (2013), who linked flooding events in Delta State to heightened waste accumulation in drains.

Table 5: Correlation coefficients for waste drivers in Warri

Variable	Correlation (r)
Population density	0.81
Industrial activity	0.74
Distance to city core	-0.62

Correlational analysis (Table 5) showed a strong positive relationship ($r = 0.81$) between population density and waste volume, while industrial activity also showed significant association ($r = 0.74$). These findings confirm the urban metabolism framework that positions waste as a product of demographic and economic flux (Kennedy et al 2011).

Table 6: Land use conflicts associated with dumpsites

Dumpsite	Adjacent Land Use	Conflict Type
Iyara	Farmland	Soil/water contamination
Uvwie/Effurun	Residential	Public health risks
NPA	River/jetty	Marine pollution

The interaction between waste volumes and planning frameworks was assessed by mapping waste against designated land uses. Table 6 highlights conflict zones where dumpsites overlap with residential, commercial, or agricultural areas, creating hazards and limiting future land use options.

Table 7: Projected waste generation in Warri

Year	Projected Volume (tons)
2024	543,541
2027	595,214
2030	704,982

Projections (Table 7) suggest that, under a 3% annual growth rate, Warri's waste could exceed 700,000 tons by 2030. Without interventions, this trend will overwhelm existing infrastructure and intensify planning conflicts.

Table 8: Planning strategies for aligning waste with urban development

Strategy	Expected Outcome
Decentralized transfer stations	Reduced transport/logistical cost
Recycling hubs in industrial zones	Resource recovery, job creation
Buffer zones around dumpsites	Minimized land use conflicts

Evidence-based strategies were distilled into planning options (Table 8). These include decentralized transfer stations, integration of recycling hubs into industrial estates, and buffer zoning around sensitive sites.

Conclusion

This study has demonstrated that the volume of waste generation in Warri is both a pressing environmental challenge and a structural determinant of urban planning outcomes. The findings reveal that annual waste generation exceeded 543,000 tons in 2024, with significant spatial and temporal variations across dumpsites. Uvwie/Effurun and NPA dominated waste flows, underscoring the strong link between industrial concentration, population density, and waste volume. Seasonal patterns, particularly the sharp peaks in the wet and festive periods, further illustrate how waste dynamics intersect with urban metabolism and strain existing infrastructure. The composition analysis revealed a predominance of biodegradable and plastic wastes, pointing to opportunities for resource recovery but also highlighting persistent risks from poorly managed residuals.

From a planning perspective, the study identified critical conflicts between waste sites and adjacent land uses, particularly residential and agricultural zones, which intensify health and ecological risks. The projection of waste growth to over 700,000 tons by 2030 signals the urgency of integrating waste realities into Warri's spatial development frameworks. Evidence-based strategies such as decentralized transfer stations, recycling hubs, and buffer zoning offer practical pathways for aligning waste management with land use planning. The implications are clear: treating waste merely as a sanitation issue is inadequate. Rather, it must be recognized as a core parameter of urban resilience, infrastructure design, and sustainable land allocation. For Warri, embedding waste considerations into planning instruments will not only mitigate environmental degradation but also strengthen the city's adaptive capacity in the face of rapid urbanization.

References

Bai X, Dawson R J, Ürge-Vorsatz D, Delgado G C, Barau A S, Dhakal S, Dodman D, Leonardsen L, Masson-Delmotte V, Roberts D C & Schultz S (2018). Six research priorities for cities and climate change. *Nature Climate Change, 12*(2), 105–112. https://doi.org/10.1038/s41558-021-01264-3

Efe S I Cheke L A & Ojoh CO (2013). Effects of Solid Waste on Urban Warming in Warri Metropolis, Nigeria. *Atmospheric and Climate Sciences*, 3(4). DOI:10.4236/acs.2013.34A002

Eneh O C (2025). *Municipal solid waste recycling, resource recovery and sustainability in Nigeria*. Springer.

Eyetan T E & Ifuwe C C (2024). "Effects of Atmospheric Pollution on Human Health in Port Harcourt: Its implication on urban planning", *Journal of Management and social Science Research*, 5 (1), 1-18. http//doi.org/10.47524/jmssr.v5i1.2

Eyetan T & Ozabor F (2021). "Oil spills deposits effect on soil physicochemical properties in Port Harcourt metropolis: Implication for agricultural planning", *Journal of Management and Social Science Research,* Vol. 2, No. 1/2, pp.45-58. *http://doi.org/10.47524/jmssr.v2i1.46*

Igere BE, Eyetan T E & Ojoh C O (2025). "Relevance of Adequate Sewage disposal on Disease Prevalence Amongst Sub-urban Communities in Delta State: A Case of Uvwie" *FUPRE Journal of Scientific and Industrial Research,* 9(2), 179-193. https://journal.fupre.edu.ng/index.php/fjsir

Kennedy C, Pincetl S & Bunje P (2011). The study of urban metabolism and its applications to urban planning and design. *Environmental Pollution, 159*(8–9), 1965–1973. https://doi.org/10.1016/j.envpol.2010.10.022

Ogu V I. & Ogbonna C (2021). Urban governance and the challenges of waste management in Nigeria. *Habitat International, 115*, 102398. https://doi.org/10.1016/j.habitatint.2021.102398

Ojoh C O, Ifuwe C C, Efetobo O & Eyetan T E. (2023). "Assessment of Industrial Waste on Soil Pollution in Asaba, Delta State, Nigeria"*, Tropical Journal of Science and Technology,* Vol. 4, No. 1, pp 37-45. http://doi.org/10.47524/tjst.v4i1.38

Onyemenam P I, Ojoh C O & Eyetan T E (2025) "Effects of Planning Laws and Policies on the development of Asaba metropolitan region in Delta State, Nigeria", *Journal of the Nigerian Institute of Town Planners, 30(1),* 75-90. Retrieved from https://nitpjournals.ng/index.php/jnitp/article/view/13

United Nations Environment Programme. (2024). *Global waste management outlook 2024*. UNEP.

World Bank. (2019). *Solid waste management: A global snapshot of challenges and opportunities*. World Bank Urban Development.

Zhang Y, Yang Q, Yu Y, & Wang T (2022). Urban metabolism: A review of current knowledge and directions for future research. *Journal of Cleaner Production, 343*, 130978. https://doi.org/10.1016/j.jclepro.2022.130978